

Studies on the release of histamine from basophils

5. Clinical evaluation

Yoshiro TANIZAKI, Haruki KOMAGOE, Michiyasu SUDO,
and Hiroshi MORINAGA

Department of Medicine, Okayama University Medical
School, Misasa Medical Branch

(Received January 7, 1985)

Introduction

Histamine release from basophils of hypersensitive individuals has been studied by using washed leucocytes (LICHTENSTEIN, L. M., et al. 1971, 1973, NORMAN, P.S., et al., 1973). These results have shown that basophils release a significant amount of histamine after stimulation with antigen and anti-IgE (LICHTENSTEIN, L. M., et al., 1970), and that the histamine release from human basophils is a valuable tool for in vitro studies of IgE-mediated, immediate hypersensitivity (RADERMECKER, M. F., et al., 1980).

During the last three years, histamine release from basophils has been extensively studied at Department of Medicine, Okayama University Medical School, Misasa Medical Branch, to clarify the pathogenesis of bronchial asthma (TANIZAKI, Y., et al., 1983b, 1983c, 1984b, 1984d, 1984f). In the following section, we summarize the results obtained for three years.

Determination of Histamine

Histamine release from basophils has been mainly examined by using washed leucocytes. Our experiments of histamine release have been carried out by using whole blood (TANIZAKI, Y., et al., 1983f, 1984a, 1984c). There are some differences between a washed leucocytes method and a whole blood method. Firstly, the incubation time of basophils with antigen and anti-IgE required to reach the maximum percent histamine

release is shorter (15 min) in the whole blood method than the time (60 min) in the washed leucocytes method (TANIZAKI, Y., et al., 1984a). Secondly, histamine release from whole blood is affected by several factors, such as total serum IgE, specific IgE antibodies and the reactivity of basophils (TANIZAKI, Y., et al. 1984e). The effect of serum factors on histamine release is more remarkable in anti-IgE-induced release than in antigen-induced release. The dose-response curves of histamine release clearly shows that the curves produced by anti-IgE are different from those by antigen. In antigen-induced histamine release, basophils release more histamine with a high concentration of antigen than with a low antigen concentration. The dose-response slope from the low to the high concentration of antigen becomes higher in accordance with higher RAST scores. On the other hand, basophils from cases with a high concentration of serum IgE require much more anti-IgE to cause maximum histamine release, because of the interaction of IgE molecules contained in their serum. In cases with a low serum IgE level (less than 300 IU/ml), a large amount of histamine was released by a low concentration of anti-IgE compared to a high concentration of anti-IgE. Thirdly, the whole blood method is easier to perform and reflects the true in vivo immediate allergic reaction better than the washed leucocytes method.

In the determination of histamine released

from leucocytes of allergic individuals, a sensitive fluorometric method has been utilized. The method is time-consuming and must be performed by a highly trained technician to get a reliable result. SIRAGANIAN, R.P., et al. (1974, 1975) has developed an automated fluorometric histamine analysis system. The method made it possible to measure histamine release more easily. Through the use of the method 30 samples per hour can be analyzed. Our histamine release experiments have been carried out by the automated fluorometric histamine analysis system. Our long-term studies on histamine release shows that the automated fluorometric histamine analysis is an easy method for the determination of extent of histamine release and that the method could be used for clinical studies of allergy (TANIZAKI, Y., et al., 1983f, 1984a).

Clinical Indications

Recent studies have clarified that phospholipid methylation caused by antigen and anti-IgE enhances membrane permeability of the target cells, inducing an increase of Ca^{2+} influx into the cells, which is accompanied by histamine release (FOREMAN, J.C., et al., 1977, HIRATA, F., et al., 1979, ISHIZAKA, T., et al., 1979, RANADIVE, N.S., et al., 1980). Our previous studies also showed that antigen causes an increase in Ca^{2+} uptake by sensitized mast cells and that inhibition of Ca^{2+} influx leads to the inhibition of histamine release (TANIZAKI, Y., et al., 1983a, 1983d, 1983e). The mechanism of histamine release through IgE receptors on the target cells seems to be similar, although MARONE, G., et al. (1981) reported that there are some differences between anti-IgE- and antigen-induced histamine release. It is important for clinical application of histamine release experiments to measure anti-IgE- and antigen-induced release from basophils of allergic individuals.

1. Basophil Reactivity to Anti-IgE

Basophil reactivity to anti-IgE can be observed by measuring histamine release from the cells. Anti-IgE-induced histamine release has been re-

ported to correlate to a certain extent with serum IgE levels (LICHTENSTEIN, L. M., et al., 1970). ASSEM, E. S. K., et al. (1981), however, showed that there was no positive correlation between the release of histamine with anti-IgE and serum IgE levels, because allergic patients' serum IgE level was very variable, but their leucocytes showed a consistently high response to anti-IgE. In our experiments, patients with a high level of serum IgE consistently showed a significant amount of histamine release when their basophils were incubated with anti-IgE. While, basophil histamine release was low in some cases and significantly high in other cases with a low serum IgE level. That is, histamine release from basophils induced by anti-IgE is variable in cases with a low serum IgE level (TANIZAKI, Y., et al., 1984d).

2. Basophil Reactivity to Antigen

Our previous studies showed that basophils participate in an attack of asthma and change morphologically after in vitro stimulation with anti-IgE and antigen (KIMURA, I., et al., 1973, 1974, 1975a, 1975b, 1981a, 1981b, 1983, In press). When basophils are incubated with antigen, histamine release and morphological changes are both observed. The morphological changes of basophils (reactive basophils) induced by antigen have a close correlation with the skin end-point titrations of antigen, RAST score and bronchial challenge tests. It is, therefore, important for detection of antigen to observe morphological changes of basophils and histamine release after addition of antigen.

1) Histamine release by house dust extract

House dust extract induces histamine release from basophils of asthmatic patients sensitive to house dust. The release of histamine induced by house dust correlates with the level of specific IgE antibodies expressed as a RAST score. The amount of histamine release induced by house dust paralleled the amount of the release caused by anti-IgE in the cases with a RAST score of 2+ or higher (TANIZAKI, Y., et al., 1984g). Skin sensitivity to house dust correlates with basophil

reactivity to the antigen. Skin sensitivity to house dust was lower in the cases with basophils low reactive to the antigen, and skin sensitivity was higher with higher reactivity of basophils. These results show that observation of histamine release from basophils is useful for detection of specific antigen and that basophil reactivity to antigen might reflect the degree of in vivo IgE-mediated allergic reaction.

2) Histamine release by *Candida albicans* extract

Candida albicans also induces histamine release from basophils of asthmatic patients sensitive to the antigen (TANIZAKI, Y., et al, 1984f, In press). Histamine release induced by *C. albicans* correlates with serum levels of specific IgE antibodies, and the amount of histamine release increases as the RAST score is higher. The release of histamine induced by *C. albicans* seems to be similar to the release by house dust. There is, however, several differences between house dust- and *C. albicans*-induced histamine release. House dust-induced release of histamine correlates with the release induced by anti-IgE in cases with a positive RAST score, but the release induced by *C. albicans* does not correlate with anti-IgE-induced release. House dust extract never causes histamine release in cases with a negative RAST score, while *C. albicans*-induced histamine release can be observed even in cases with a negative RAST score. These results shows that non-IgE-mediated reaction participates in *C. albicans*-induced release of histamine.

References

1. ASSEM, E.S.K. and ATTALLAH, N. A. (1981): Increased release of histamine by anti-IgE from leucocytes of asthmatic patients and possible heterogeneity of IgE. *Clin. Allergy* **11**, 367-374.
2. FOREMAN, J.C., HALLETT, M.B. and MONGER, J.L. (1977): The relationship between histamine secretion and ⁴⁵Calcium uptake by mast cells. *J. Physiol.* **271**, 193-214.
3. HIRATA, F., AXELROD, J. and CREWS, F. T. (1979): Concanavalin A stimulates phospholipid methylation and phosphatidylserine decarboxylation in rat mast cells. *Proc. Natl. Acad. Sci. USA* **73**, 4813-4816.
4. ISHIZAKA, T., FOREMAN, J.C., STERK, A. R. and ISHIZAKA, K. (1979): Induction of calcium uptake across the rat mast cell membrane by bridging IgE receptors. *Proc. Natl. Acad. Sci. USA* **76**, 5858-5862.
5. KIMURA, I., MORITANI, Y. and TANIZAKI, Y. (1973): Basophils in bronchial asthma with reference to reagin-type allergy. *Clin. Allergy* **3**, 195-201.
6. KIMURA, I., TANIZAKI, Y., TAKAHASHI, K., SAITO, K., UEDA, N. and SATO, S. (1974): Emergence of basophils at the site of local allergic reaction using a skin vesicle test. *Clin. Allergy*, **4**, 281-285.
7. KIMURA, I., TANIZAKI, Y., SAITO, K., TAKAHASHI, K., UEDA, N. and SATO, S. (1975a): Appearance of basophils in the sputum of patients with bronchial asthma. *Clin. Allergy* **5**, 95-101.
8. KIMURA, I., TANIZAKI, Y., SATO, S., TAKAHASHI, K., SAITO, S. and UEDA, N. (1975b): Morphological changes of basophils in immunological reactions-effect of sodium cromoglycate. *Clin. Allergy* **5**, 181-187.
9. KIMURA, I., TANIZAKI, Y., SATO, S. and TAKAHASHI, K. (1981): Differences in response to anti-IgE and to anti-IgG in basophils from patients with bronchial asthma. *Clin. Allergy* **11**, 31-36.
10. KIMURA, I., TANIZAKI, Y., SATO, S., TAKAHASHI, K., SAITO, S. and UEDA, N. (1981b): Supravital observation of in vitro basophils in immunological reactions. *Clin. Allergy* **11**, 37-41.
11. KIMURA, I., TANIZAKI, Y., GODA, Y., SASAKI, Y., AKAGI, K., KOBAYASHI, M. and SODA, R. (1983): New in vitro method for detecting asthma allergen. Counting reactive basophils after addition of allergen. *Clin. Allergy* **13**, 99-105.
12. KIMURA, I., TANIZAKI, Y., GODA, Y.,

- KOMAGOE, H. and KITANI, H. (In press): Decrease in reactivity of basophils by immunotherapy with house dust extract. *Clin. Allergy* 15.
13. LICHTENSTEIN, L. M., NORMAN, P. S. and WINKENWERDEN, W. L. (1971): A single year of immunotherapy for ragweed hay fever. Immunologic and clinical studies. *Ann. Intern. Med.* **75**, 663-671.
 14. LICHTENSTEIN, L. M., ISHIZAKA, K., NORMAN, P. S., KARGEY-SOBOTKA, A. and MILL, B. M. (1973): IgE antibody measurements in ragweed hay fever. Relationship to clinical severity and the results of immunotherapy. *J. Clin. Invest.* **52**, 472-482.
 15. LICHTENSTEIN, L. M., LEVY, D. A. and ISHIZAKA, K. (1970): In vitro reversed anaphylaxis: characteristics of anti-IgE mediated histamine release. *Immunology* **19**, 831-842.
 16. MARONE, G., KAGEY-SOBOTKA, A. and LICHTENSTEIN, L. M. (1981): IgE-mediated histamine release from human basophils: Differences between antigen E- and anti-IgE-induced secretion. *Int. Archs Allergy Appl. Immunol.* **65**, 339-348.
 17. NORMAN, P. S., LICHTENSTEIN, L. M. and ISHIZAKA, K. (1973): Diagnostic tests in ragweed hay fever. *J. Allergy Clin. Immunol.* **52**, 210-224.
 18. TANIZAKI, Y. and TOWNLEY, R. G. (1983a): Effect of BSA on Ca^{2+} influx in mast cells stimulated by ovalbumin. *Int. Archs Allergy Appl. Immunol.* **70**, 143-145.
 19. TANIZAKI, Y., KOMAGOE, H., SUDO, M., et al. (1983b): Studies on the release of histamine from basophils. 1. Determination of histamine from whole blood by an automated fluorometric histamine analysis system. The Papers of the Institute for Thermal Spring Research, Okayama University.
 20. TANIZAKI, Y., KOMAGOE, H., SUDO, M., et al. (1983c): Studies on the release of histamine from basophils. 2. Histamine release induced by house dust and anti-IgE. The Papers of Institute for Thermal Spring Research, Okayama University.
 21. TANIZAKI, Y., KOMAGOE, H., SUDO, M., et al. (1983d): Inhibitory effect of the Ca^{2+} antagonist nifedipine on histamine release from rat peritoneal mast cells. *Acta Med. Okayama* **37**, 207-211.
 22. TANIZAKI, Y., AKAGI, K., LEE, K. N. and TOWNLEY, R. G. (1983e): Inhibitory effect of nifedipine and cromolyn sodium on skin reactions and ^{45}Ca uptake and histamine release in rat mast cells induced by various stimulating agents. *Int. Archs Allergy Appl. Immunol.* **72**, 102-109.
 23. TANIZAKI, Y., KOMAGOE, H., SUDO, M., MORINAGA, H., et al. (1983f): IgE-mediated histamine release from whole blood in atopic asthmatics. *Jpn J. Allergol.* **32**, 1079-1083.
 24. TANIZAKI, Y., KOMAGOE, H., MORINAGA, H., KITANI, H., GODA, Y. and KIMURA, I. (1984a): Allergen- and anti-IgE-induced histamine release from whole blood. *Int. Archs Allergy Appl. Immunol.* **73**, 141-145.
 25. TANIZAKI, Y., KOMAGOE, H., SUDO, M., et al. (1983b): Studies on the release of histamine from basophils. 3. Correlation between basophil reactivity to anti-IgE and blood eosinophilia. The Papers of Institute for Thermal Spring Research, Okayama University **54**, 35-38.
 26. TANIZAKI, Y., KOMAGOE, H., SUDO, M., MORINAGA, H., et al. (1984c): Histamine release from whole blood induced by anti-IgE: relationship to patient age, age at onset and serum IgE levels. *Acta Med. Okayama* **38**, 275-280.
 27. TANIZAKI, Y., KOMAGOE, H., SUDO, M., KITANI, H., NAKAGAWA, S. and KIMURA, I. (1983d): Basophil reactivity to anti-IgE and allergen in asthmatic subjects: relation to house dust allergy. *Jpn J. Clin. Immun.* **7**, 177-181.
 28. TANIZAKI, Y., KOMAGOE, H., SUDO, M., MORINAGA, H., et al. (1984e): Effect of a serum factor on IgE-mediated histamine release from whole blood. *Acta Med.*

Okayama 38, 381-387.

29. TANIZAKI, Y., KOMAGOE, H., SUDO, M., MORINAGA, H., et al. (1984f) : Studies on the release of histamine from basophils. 4. Difference between house dust- and Candida-induced secretion. The Papers of The Institute for Thermal Spring Research, Okayama University 55, 7-10.
30. TANIZAKI, Y., KOMAGOE, H., SUDO, M., MORINAGA, H., et al. (1984g) : Reactivity of sensitized human basophils as expressed by histamine release. Jpn J. Allergol. 33, 463-467.
31. TANIZAKI, Y., KOMAGOE, H., SUDO, M., KITANI, H., NAKAGAWA, S., et al. (In press) : Participation of specific IgG antibodies in basophil histamine release induced by Candida albicans. Jpn J. Allergol.
32. RADERMECKER, M.F. (1980) : Allergen-induced histamine release from whole blood. Clinical evaluation. Int. Archs Allergy Appl. Immunol. 63, 415-423.
33. RANADIVE, N. S. and DHANARI, N. (1980) : Movement of calcium ions and release of histamine from rat mast cells. Int. Archs Allergy Appl. Immunol. 61, 9-18.
34. SIRAGANIAN, R. P. (1974) : An automated continuous flow system for the extraction and fluorometric analysis of histamine. Analyt. Biochem. 57, 383-394.
35. SIRAGANIAN, R.P. (1975) : Refinements in the automated fluorometric analysis system. J. Immunol. Method 7, 283-290.

好塩基球からのヒスタミン遊離に関する研究.

5. 臨床的評価

谷崎勝朗, 駒越春樹, 周藤真康, 森永 寛

岡山大学三朝分院内科

過去3年間にわたり, 気管支喘息患者末梢血好塩基球からのヒスタミン遊離について, ヒスタミン自動分析装置を用いて全血法により検討を加えてきた.

1. 抗ヒト IgE によるヒスタミン遊離は, 症例間で著しい差がみられた. この際血清 IgE 値が高値を示す症例では全般的に高度なヒスタミン遊離がみられたが, 血清 IgE 値が正常かまたはむしろ低値を示す症例のヒスタミン遊離は, かなり高度なものから全くみられないものまでさまざまであった.
2. ハウスダストやカンジダなどの抗原物質によるヒスタミン遊離は, 特異的 IgE 抗体依存性であり, 抗体価が上昇するにつれヒスタミン遊離は高度となる傾向がみられた. しかし, ハウスダストとカンジダによるヒスタミン遊離には若干の差がみられた. すなわち, ハウスダストと抗ヒト IgE によるヒスタミン遊離の間には密接な関連がみられたが, カンジダと抗ヒト IgE の間には全く関連がみられなかった. またカンジダによるヒスタミン遊離においては, IgE系反応以外の反応が関与する可能性が一部示唆された.